

## American Peony Society Cultivar Registration Form (EXPLAINED)

PROPOSED NAME: Proposed name for new cultivar Date: Month/Day/Year  A cultivar is defined by the ICNCP as "an assemblage of plants that has been selected for a particul combination of attributes and that is clearly distinct, uniform, and stable in these characteristics as propagated by appropriate means retains those characteristics." This means that a seedling, no madesirable, is NOT a cultivar until it has been propagated and shown to be stable in those desirable not qualify as a cultivar, it should not be formally named or registered.	nd that when atter how good or
Name derivation <u>What is the origin or meaning of the name.</u> In some cases this will be obvious while in other cases explanation may be needed. If named for a precently deceased, permission must be obtained from that person (or heirs) before the name can be	
ORIGINATOR:The person who raised the peony Name / Address	Name Street
REGISTRANT (if different from originator): The person registering the peony. Name / Address	City, State, Zip Code or City, Province-Region, Postal Code, Country
SEEDLING NUMBER or GARDEN NAME: Number, code, or garden name by which seedling registration  This is particularly important for peonies which were distributed or exhibited prior to formal name	
registration.  FIRST YEAR BLOOMED:YEAR_  This is important to gauge the maturity of the plant with respect to flowering and flower form stable.	ility.
FIRST YEAR PROPAGATED:YEAR	s of a plant, and al seedling with the
If Yes, explain organization giving patent or trademark, patent number, etc  Has cultivar been exhibited and/or received awards? Yes No check one	check one
If Yes, explain give number or name under which it was shown and who (Organization) gave t	the award
PARENTAGE: Pod ParentPlant name or number which produced the seed Pollen ParentPollen parent name or number (source) Unknown	
Unknown In some instances the parentage from both sides will be unknown. Check if applicable.  TYPE: (Cultivar Group):	
Lactiflora Group check if applicable  These are the common garden peonies descended from the Chinese herbaceous peonies. They are di chromosome number and there are more of them than any other group. Crosses between members on not considered hybrids. Multiple blooms per stem almost always occur in this group. As a group, the bloom. Also in this group are the "Lactifloras of hybrid origin" which result from hybrids being cros to the point they become visually indistinguishable from them.	of this group are ey are latest to
Herbaceous Hybrid check if applicable  These are herbaceous peonies which have more than one species in their parentage. Some have ped four or more species in their makeup. Typically one bloom per stem, but sometimes more. Colors temore vibrant than in the lactifloras.	
Itoh Group check if applicable  Itoh Group peonies are hybrids having Lactiflora Group peonies as seed parent and Lutea Hybrid C  pollen parent. The term "Intersectional" applies to crosses between two different sections of the gen  general descriptive term. Itoh applies to a specific cross as noted above. Features of most of these to  like foliage and flowers, with herbaceous plant habit.	us and is thus a

Suffruticosa Group check if applicable
The group includes the traditional Chinese and Japanese tree peonies, any of the purported Rockii Hybrids, and any
other cultivars resulting from crosses within the subsection Vaginatae. Subsection Vaginatae includes all woody species
except for P. delavayi (lutea) and P. ludlowii.
Lutea Hybrid Group Peony check if applicable
These are woody peony cultivars originally derived by crossing between members of the subsection Delavayanae (P.
delavayi (lutea) and P. ludlowii) as one parent, and members of the Suffruticosa Group as the other parent. Crosses
within the Lutea Hybrids Group are also included.
Species Selection check if applicable
Examples may be selections from wild populations which show characteristics desired for introduction into general
cultivation.
Other check if applicable
For registration purposes, here would be listed any other peony that does not fit any of the above descriptions.
Examples may be selections from hybrids between species within the Delavayi group could be listed here.

SEASON OF BLOOM: Very Early	Early   Early/Mid	Midseason	Mid/Late [ ]Lat	e Very Late
-----------------------------	-------------------	-----------	-----------------	-------------

	Bloom Season Guide Tab	
Complete list may be	e found on the Internet at http:	://bloomdate.paeonia.com/
SEASON MIDPOINT	BLOOM SEASON	SEASON BOUNDARY
	1 1 1 7	← P. tenuifolia (single)
Nosegay →	Very Early	
77 / 1	20 Mar. 14	← Windchimes
Laddie →	Early	
		← Picotee
Dawn Glow →	Early-Mid	
	277	← Archangel
RED CHARM $\rightarrow$	Midseason	
		← Many Happy Returns
Kay Tischler →	Mid-Late	
	1	← Mrs. Livingston Farrand
Isani Gidui →	Late	
		← A. B. Franklin
Myrtle Gentry →	Very Late	
/ × /		← Vivid Rose

In the representation above, the column on the right lists cultivars that mark the approximate boundaries of bloom interval. Those on the left mark the mid-point of bloom interval. It is not invariably precise, but it will serve as a useful guide, as explained in the following paragraphs.

Bloom data has been collected for almost 1000 different peony cultivars, including species and a few tree peonies, and for most, in multiple years (see http://bloomdate.paeonia.com/). When the data is analyzed, it shows that there can be more than seven weeks of peony bloom of one sort or another. Weather plays a large part in expanding the season in cooler years, or compressing it during warm years.

There are almost as many days in which peonies can be expected to bloom before Red Charm, as there are ones in which they bloom after Red Charm, and so Red Charm makes a good starting point. With Red Charm as the mid-point, it suggests an odd number of periods, and this works out very well too with naming these bloom periods. We can make seven bloom periods, each of 6 to 8 days duration depending on the weather in any particular year.

This is not a precise system, there being so many variables from one garden to the next, as well as differences in the micro-climates within those gardens, but in general it does make a much better system for describing when a

particular peony can be expected to bloom than anything we've had before. Access to the table containing all data will still be useful for the gardener wishing to expand their peony bloom season, but the seven letter designations will give a broad idea of when something might be expected to bloom.

This is not a new idea, it's just that it took 40 years since William Krekler proposed such a scheme in 1960 before there were people willing to do some detailed data collection and analysis. Krekler had only 5 divisions (Very Early, Early, Mid, Late, Very Late) and maybe these are a better option than seven. His suggestion was that peony bloom designations should begin to have some basis in reality, and reflect the true bloom season of ALL peonies, not just the lactifloras.

## **FLOWER**

Flowers change as they age. For descriptive purposes it is assumed that observations will be made on a fresh bloom. In certain types, colors may progress through a series of changes which add to garden effect, or contribute to diagnostic tools, and these should be mentioned.

DТ	$\cap$	X7	$\mathbf{r}\mathbf{p}$	FO	١D٨	л.
ГI.	ハノ	vv	$\Gamma_{I}\Gamma_{I}$	FU	יותוי	/I ·

LIO M LIL LOIMI.
Single - check if applicable. Typically resembles the wild form with 5 petals, pollen bearing stamens, and functional carpels. Singles in cultivation may have up to 15 petals but still be classed as singles if the petals are spread in a saucer shape rather than cup shaped, ruffled, or otherwise give the impression of bulk. Japanese - check if applicable. Diagnostic of Japanese flower form are staminodes. These are abortive or transformed stamens in which that origin is still recognizable. Edges of the staminodes are thickened and often contain viable pollen, but it is encased in tissue and not otherwise available. The original peonies of this form were imported from Japan where the lack of pollen drop was a desirable feature. Anemone - check if applicable. Stamen transformation in the anemone form has progressed to the point where all visible evidence of stamen origin, except for yellow color, has disappeared. These structures are termed petalodes and resemble petals. Some of these petalodes are very narrow, and yellow, giving the effect of a Japanese form peony. Other petalodes are wider and as size increases, the flowers begin to approach the bomb flower form. Semi-Double - check if applicable. Quite a variety of forms are included in this designation. They all have in common prominent stamens, and a bulking of the petalage, either through partial transformation of stamens into petalodes, increased number of guard petals, or guard petal structure which adds visual bulk to the flower. Bomb - check if applicable. Petalodes have progressed in size to almost the same size as the guard petals, and are almost always the same color as the guards. Overall effect is a ball sitting on a plate, the plate being formed of the larger guard petals at the base of the flower, and the shorter central petals forming the ball. Red Charm and Mons. Jules Elie are examples. Full Double - check if applicable. In the ultimate expression of this form, all stamens and carpels are transformed into petals. Scattered st
Other (explain) - check if applicable. if it doesn't seem to fit any of the above classifications, describe flower form
here. Not all peonies cooperate by falling into easily defined flower form, and some regularly show more than one form on the same plant.
FLOWERS PER STEM (average number): Includes the main flower bud and any side buds.
FLOWER SIZE: Size of the main flower. Generally the diameter is specified, but the height of the bloom could be recorded too.
FLOWER PRESENTATION:Up facingOut FacingDown Facing  Almost every cultivar will have up facing flowers. Out Facing flowers are held at more or less right angles to the stem, such as in Windchimes and Saunders' Early and Late Windflowers. Down Facing flowers are those where the bloom hangs down in a position not suitable for normal viewing. This is typically found in some tree peonies, Souvenir de Maxime Cornu being a prime example.
FRAGRANCE:Very Fragrant Some Fragrance No FragranceObjectionable Odor Many peonies have some fragrance, but not all have a strong aroma. A few have an objectionable odour.
PETAL COLOR: Primary: The main petal color. RHS If available, a color chart # can be used, but specify the authority. In this case, RHS signifies Royal Horticultural Society.  Secondary: Edging, streaking, overlay, etc RHS  COLOR PATTERNS: If flares or blotches are present, or if petals are of more than one distinct color, or a blend of
colors, describe color pattern.

FLARE	S: _Yes	No.	Color				RHS	
Margin o	of flare:	Sharp	edge	Blurred edge Blee	eding edge			
				length % or frac				
				s of intense pigmentat				
								but are also found in a few
								on the reverse side of petals in
								e flares found on the front side
				led obverse blotches. ey seldom contribute				y coloured than the
	surround	ing area	a anu m	ey seldom contribute	much in the way	y or beauty	y.	
GUARI	) PETAL	FORM	M (selec	t all that apply):				
Flat	Cupp	oed	Twisted	dRoundedPo	ointedRuff	ledFr	rilledNotched	
	er (expla							
	e width o							
				age Number. Color_			RHS	
Hairine	ss: S	mooth	Spa	arse Moderate _	Very Hairy			
				RHS			Feathered	
STAME	NS: _Y	esN	o POLL	LEN: _Yes _No SE	EDS: <u>Yes</u>	_No		
STAME	N LENC	ΉΤH:	None	Less than ½"	½" to ¾"	More th	ian ¾"	
FILAM	ENT CO	LOR: I	Base				_ RHS	
Upper_					RHS			
								ts. The filament is sometimes
		n one co	olor, witl	n the lower portion (ba	ase) often darke	r than the	e upper half. "Uppe	r" here does <u>not</u> mean the
стамт	anther. NODES:	Voc	No					
	NODES. NODE S							
				nost normal, therefore	alagaler wagamah	lo atomony		
	ar – narı				closely resemb	ie stameni	s.	
				er sides. htly wider towards the	a tim			
	ulate – s			itty wider towards til	e up.			
_	$\operatorname{er} - \operatorname{Desc}$							
STAMI	NODE C	OI OR	AND C	COLOR PATTERNS	NCSLA-800			
	Staminod diagnosti they don' staminod	des are posterior description	partially re. A rule exposed p same as	transformed stamens e of thumb is that star pollen. There is usual	s seen in some fl minodes have vi ly a yellow edge ses there is ofte	sual evide or tip, the n a differe	ence of their transfo ough sometimes the ence in texture betw	e flower form they are a cormation from stamens, but e color of the body of the ween the edges and centre of narrow.
DISK:								
	of Devel	opmen	t: _ W	ell Developed and C	Complete A	Few Pro	jections Obscu	are.
Color:							~DA	
	In herbac flower. The ring arou	his is thand the	ne disk, s base of t	sometimes referred to	as the staminod	dal disk. A	well developed dis	ne carpels at the base of the sk forms an almost complete ounded projections, or
SHEAT				l Complete				
Color _	111	one	_ 1 01 010	complete	RHS			
	form the	sheath,	a paper		ies and herbace he carpels. The	sheath ma		extremely well developed to ring degrees of development
					PLANT			
LEAFL	ETS:	Finely	Cut	_Broadly CutNa	arrow Pointed	Broad	d PointedBroa	ad Blunt Irregular
LEAF (	COLOR C	ON EM	ERGEN	NCE FROM GROUN	ND:			

LEAF COLOR WHEN MATURE:GreenYellow-GreenBlue Green
Secondary Color: Pink Purple
Other (explain)
Stems and foliage often emerge from the ground in shades of red or purple, and then become green as the season progresses. Some retain red stems, or partially so, for varying lengths of time. These characteristics can be both diagnostic features and seen as desirable for plant and garden interest. Secondary leaf colour may refer to variegation, the underside of the leaf, the purplish cast seen on the foliage of some woody peonies, or any other such thing as may be notable.
AVERAGE HEIGHT AT MATURITY: DOES PLANT NEED SUPPORT? Yes No
GROWTH HABIT:Low/BroadSpreadingUprightNarrow Upright.

## WRITTEN DESCRIPTION

A description written by the applicant is always useful, and this is the place to include information not asked for in the form. Please provide a description of the cultivar that distinguishes it from all others of a similar type and color, and any other information you feel is important or potentially of interest.

## **PHOTOGRAPHS**

Include with your request for registration a color photo, color slide, or color digital image of the flower and of the foliage. A single photograph clearly showing details of flower and foliage is acceptable, though three photos, flower, foliage, and entire plant, provide the best record. The photo of the foliage can be either a single complete stem (herbaceous) or a leaf (woody or intersectional).